

UNIVERSAL SCALING OF THE 3:2 TWIN-PEAK QUASI-PERIODIC OSCILLATION FREQUENCIES WITH BLACK HOLE MASS AND SPIN REVISITED

XIN-LIN ZHOU¹, WEIMIN YUAN¹, HAI-WU PAN¹, ZHU LIU¹

¹ National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing, 100012, China; zhouxl@nao.cas.cn

Draft version December 16, 2014

ABSTRACT

We discuss further observational support of an idea formulated a decade ago by Abramowicz, Kluźniak, McClintock and Remillard. They demonstrated that the 3:2 pairs of frequencies of the twin-peak black hole (BH) high-frequency quasi-periodic oscillations (QPOs) scale inversely with the BH masses and that the scaling covers the entire range from stellar to supermassive BHs. For this reason, they believed that the QPOs may be used for accurate measurements of masses and spins of BHs.

Subject headings: black hole physics – X-rays: galaxies – X-rays: binaries – galaxies: general

1. INTRODUCTION

Multiple quasi-periodic oscillations (QPOs) are often found in the power spectrum of X-ray variability in X-ray binaries. High-frequency QPOs (HFQPOs; 40 – 450 Hz) have been detected in a few X-ray binaries (Remillard 2004). These HFQPOs may correspond to the frequency at the innermost stable circular orbit for Schwarzschild black hole (BH) with masses of 5 – 15 $M_{\odot}$. These QPO frequencies do not change significantly despite sizable changes in the X-ray luminosity. This suggests that the HFQPO frequencies are primarily dependent on the mass and spin of the BH (Remillard & McClintock 2006). Thus HFQPOs are likely to offer the most reliable measurement of spin once the BH mass is known and correct model is applied (e.g., Abramowicz & Kluźniak 2001; Rezzolla et al. 2003).

QPOs have been detected in a few objects, e.g., the ultra-luminous X-ray sources NGC 5408 X-1 (Strohmayer et al. 2007), M82 X-1 (Strohmayer & Mushotzky 2003; Feng & Kaaret 2007; Caballero-García et al. 2013; Pasham et al. 2014), the Seyfert galaxy RE J1034+396 (Gierliński et al. 2008; Alston et al. 2014a), and the tidal disruption event Swift J1644+57 (Reis et al. 2012). Are these QPOs the analogies of HFQPOs? Can the BH spin play a role? Here we study the correlation between the QPO frequency and BH mass for these objects. We find a universal scaling relation from Galactic X-ray binaries to active galactic nuclei.

There is a simple theoretical reason for the inverse mass scaling of the BH QPOs frequencies. Indeed, strong gravity has a characteristic lengthscale $R_G = 2GM/c^2$, which formally corresponds to a characteristic frequency $\nu_G = c/R_G = c^3/(2GM) \sim 1/M$. If the observed BH QPO frequencies reflect some orbital (geodesic) frequencies, then one would expect that $\nu_{\text{QPO}} = [c^3/(2GM)]f(x, a)$, where $x = r/R_G$ is a dimensionless radius at which the oscillation occurs, and a is the dimensionless spin. Abramowicz & Kluźniak (2001) were the first to noticed in the observational data — that the observed twin-

peak QPO frequencies are in a 3:2 ratio,¹ and argued that the coherence of QPO, discussed above, suggests a *resonance*. In this case, $x = x_{\text{res}}(a)$ is fixed by the very nature of the resonant oscillations that are behind the observed 3:2 ratio of the twin-peak QPO frequencies, and $\nu_{\text{QPO}} = [c^3/(2GM)]f(x_{\text{res}}(a), a) = [c^3/(2GM)]F(a)$, with $F(a)$ being known for any specific type of resonance. Thus, the “3:2 QPO frequencies” depend on mass and spin through a known relation, $\nu_{\text{QPO}} = \nu_{\text{QPO}}(M, a)$, and measuring ν_{QPO} may be an accurate tool for measuring mass and spin. This idea, first applied by Abramowicz and Kluźniak (2001) for the Galactic BH QPOs was later employed by numerous authors, e.g. Török et al. (2005), also for SgrA* Török (2005), and more recently by Abramowicz & Liu (2012) for the Swift J164449.3+573451 supermassive BH.

Our Letter is based on the paper by Abramowicz et al. (2004). They stressed the importance of the universal $\nu \sim 1/M$ scaling and suggested that it may be used for a definite mass measurement of the hypothetical intermediate BH, if the twin-peak 3:2 QPOs would be discovered for such sources. Indeed, Pasham et al. (2014) have recently discovered a 3:2 twin-peak QPO in the ultra-luminous source M82 and claimed, following the Abramowicz et al. (2004) suggestion, that this is a 400 $M_{\odot}$ BH. Throughout this work, we assume the concordance cosmology of $h_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$, and $\Omega_{\Lambda} = 0.73$.

2. OBJECTS WITH (HF)QPO DETECTION

Table 1 lists the objects with the (HF)QPO detections. Four BH X-ray binaries, GRO J1655 – 40, XTE J1550 – 64, GRS 1915 + 105, and H 1743 – 322 display a pair of HFQPOs with a 3:2 frequency ratio. The frequen-

¹ Titarchuk (2002) wrote about the chronology of these observational discoveries: “Recently Abramowicz et al. (2003) drew the attention of the community to the possible presence of the 3:2 resonance in QPOs for the neutron star sources. A similar effect was also pointed out earlier by Abramowicz & Kluźniak (2001) and Remillard et al. (2002) for the BH sources.” Note also that neither in Strohmayer (2001a), nor in Strohmayer (2001b), the ratio 3:2 is mentioned. Although Strohmayer discovered the twin-peak BH QPOs, he did not notice their 3:2 frequency ratio.

cies of X-ray binaries are listed in Table 1 for the stronger QPO that represent $2 \times \nu_0$. The former three X-ray binaries have well-measured dynamical BH mass. Literatures by different authors gave slightly different values for BH mass. Recently, Reid et al. (2014) measured a trigonometric parallax for the X-ray binary GRS 1915 + 105. This gives a direct distance estimate and a revised estimate for the BH mass of $12.4^{+2.0}_{-1.8} M_\odot$. This is still in the range of $10 - 18 M_\odot$ used in Remillard & McClintock (2006).

QPOs have been detected in the ultra-luminous X-ray source M82 X-1 (Strohmayer & Mushotzky 2003; Feng & Kaaret 2007). Caballero-García et al. (2013) reported on the detection of (46 ± 2) mHz QPOs in the power density spectra of M82 X-1. A comparison of the frequency of these QPOs with previous detections supports the 3:2:1 frequency distribution. If it is true, this implies a BH mass of $\sim 10^4 M_\odot$ for M82 X-1. However, Pasham et al. (2014) presented twin-peak (3:2 frequency ratio) X-ray QPOs from M82 X-1 at frequencies of 3.32 ± 0.06 Hz and 5.07 ± 0.06 Hz. They also estimated a BH mass of $415 \pm 63 M_\odot$ using the relativistic precession model.

The ultra-luminous X-ray source NGC 5408 X-1 shows a QPO at 20 mHz (Strohmayer et al. 2007). They also found evidence for a second QPO peak at 15 mHz. The frequency ratio of this QPO pair is inconsistent with 3:2 but is consistent with a 4:3 ratio. The BH mass of this source is still under debate. Recently, Huang et al. (2013) presented X-ray timing and spectral analysis to support a large BH mass of $\sim 10^5 M_\odot$. The X-ray light curve of one 90 ks XMM-Newton observation of the Seyfert galaxy RE J1034 + 396 shows an ~ 1 hour X-ray periodicity (Gierliński et al. 2008). The QPO feature is detected in five low-flux/spectrally harder observations (Alston et al. 2014a). This source is a narrow line Seyfert 1 galaxy. The BH mass of $4^{+3}_{-1.5} \times 10^6 M_\odot$ for this object is derived from X-ray variability amplitude (Zhou et al. 2010). This is broadly in agreement with the value of $(1 - 4) \times 10^6 M_\odot$ derived from its bulge stellar velocity dispersion (Bian & Huang 2010).

Swift source J164449.3 + 573451 (Swift J1644 + 57) at redshift $z = 0.3543$ recently experienced a powerful outburst, most likely caused by a tidal disruption of a star by the massive BH at the center of the source. During the recent X-ray outburst, the source was similar to a small-scale blazar (Bloom et al. 2011), with a relativistic jet (Burrows et al. 2011). Reis et al. (2012) detected a firm QPO from this object with a centroid frequency ~ 4.8 mHz. It has been suggested to be an intermediate-mass BH in Swift J1644 + 57 (Abramowicz & Liu 2012; Shen & Matzner 2014). Miller & Gültekin (2011) combined X-ray and radio observations of Swift J1644 + 57 to constrain BH mass of $\sim (3.16^{+35.6}_{-2.9}) \times 10^5 M_\odot$.

The QPOs have been detected from the infrared (Genzel et al. 2003), X-ray (Aschenbach et al. 2004a) and radio bands (Miyoshi et al. 2011) in the Galactic Center Sgr A*. X-ray observations have revealed a 3:2:1 frequency ratio for Sgr A* (Aschenbach et al. 2004a), although the significance of the QPOs is still questionable. The dense gas cloud called G2 moving towards Sgr A* (Gillessen et al. 2012) will produce strong flare at multiple bands and will test the possible QPO emission in Sgr A*.

3. RESULTS AND DISCUSSION

Figure 1 shows the relationship between the QPO frequency and the BH mass listed in Table 1. Three BH X-ray binaries, XTE J1550–564, GRO J1655–40, and GRS 1915 + 105, have well-measured dynamical BH masses. The HFQPO frequencies from the three BH X-ray binaries appear to scale inversely with mass. The relationship based on these three stellar-mass BHs for the stronger $2 \times \nu_0$ frequency is (Remillard & McClintock 2006),

$$\nu(Hz) = 1862(M_{\text{BH}}/M_\odot)^{-1}. \quad (1)$$

We call this relationship the universal scaling relation. This relation is plotted in Figure 1 as the solid line, and extrapolated to the higher-mass range. If these HFQPOs are indeed general relativity oscillations, then the simple scaling with BH mass may suggest that three stellar BHs have similar values of the dimensionless spin parameter, a (Remillard & McClintock 2006). The spins for the three BHs have been measured from the X-ray continuum fitting (McClintock et al. 2011) and X-ray reflection method (Reynolds 2014). GRS 1915 + 105 (McClintock et al. 2006; Blum et al. 2009) has an extremely high spin value ($a \sim 0.99$), while GRO J1655–40 (Reis et al. 2009) and XTE J1550–564 (Steiner et al. 2011) has a moderate spin value.

It is suggested that an HFQPO with the harmonic pairs of frequencies in a 3 : 2 or 3 : 1 ratio arises from some type of resonance mechanism (Abramowicz & Kluźniak 2001; Remillard et al. 2002; Abramowicz et al. 2004). The results from the HFQPOs using resonance model (Abramowicz & Kluźniak 2001) or another model (Rezzolla et al. 2003) also suggest moderate or high spin values for the three stellar BHs.

Based on the resonance model, for a given spin value, the correlation between HFQPO frequency and BH mass is determined. The relationship derived from a model of 3 : 2 resonance and $a = 0.996$ is (Aschenbach 2004b),

$$\nu(Hz) = 2030.8(M_{\text{BH}}/M_\odot)^{-1}. \quad (2)$$

As a comparison, we also show the relationship derived from a model of 3 : 1 resonance and $a = 0.996$ (Aschenbach 2004b),

$$\nu(Hz) = 3068.9(M_{\text{BH}}/M_\odot)^{-1}. \quad (3)$$

As shown in Figure 1, they are quite close to the universal scaling relation.

Swift J1644+57 is a tidal disruption of a star by a dormant BH. The BH mass estimated from radio observations (Miller & Gültekin 2011) for this object well follows the universal scaling relation, although with a large error bar (see Figure 1). Swift J1644+57 shows evidence for the relativistic jet activity (Shao et al. 2011), a probable two-component jet (Liu et al. 2012). The 4.8 mHz QPO could be associated with a second, narrower jet periodically sweeping the observer line of sight (Wang et al. 2014). If the 4.8 mHz QPO is due to the same mechanism as what produces the HFQPOs in Galactic X-ray binaries, a very realistic possibility is that the BH in Swift J1644+57 has a mass closer to the most likely mass of $3 \times 10^5 M_\odot$ (Abramowicz & Liu 2012), and has a high spin (Wang et al. 2014). The rapid spin of a BH may be a necessary condition to generate a relativistic jet.

It is very interesting that a few objects well follow the universal scaling relation. Note that RE J1034+396 is a Seyfert galaxy, NGC 5408 X-1 and M82 X-1 are ultra-luminous X-ray sources, and Swift J1644+57 is a tidal disruption of a star by a dormant BH. They are different types of BHs and have different formation histories. All these BHs follow the universal scaling relation defined from the three Galactic X-ray binaries. This suggests that these BHs have similar spin values, which are very likely moderate or high spin values.

The spins of about a dozen of stellar BHs and about 20 supermassive BHs have been measured from the X-ray reflection method (Reynolds 2014). Many objects have moderate or high spin parameters. This implies that the cosmic processes to spin up BHs are very effective. Kerr BHs may be common at all scales (Volonteri et al. 2005). The narrow-Line Seyfert 1 galaxy RE J1043+396 slightly deviates from the universal scaling relation. If the BH mass of RE J1034+396 is close to the most likely mass of $4 \times 10^6 M_\odot$ (Zhou et al. 2010), it may have a lower spin than the stellar mass BHs. This is in good agreement with the average spin measured from the composite X-ray spectrum of narrow-line Seyfert 1 galaxies (Liu et al. 2014). If the QPOs detected in Sgr A* are real, the well measured dynamical BH mass of Sgr A* (Gillessen et al. 2009) follows the 3:1 resonance line (see Figure 1). The BH in Sgr A* may have an extreme spin parameter (Aschenbach 2004b).

Stellar BHs, such as BH X-ray binaries and micro-quasars, form via stellar processes. Accretion and angular momentum extraction cannot significantly change the spin parameter of a BH in a binary system (King & Kolb 1999; Zhang 2013). Thus the spin parameter of a stellar BH reflects the value at formation. For massive BHs, recent studies link the evolution of BH spin to

the distribution of orientations of episodic accretion periods (e.g., Wang et al. 2009). Most massive BHs would spin down if the orientations of the accretion disks are isotropic; most massive BHs would spin up if the distribution of the orientations is highly anisotropic or if accretion is coherent for a long time (Dotti et al. 2013). The formation history of intermediate-mass BH remains unknown. It is likely that they are relic BHs in the early universe, with the spin obtained from the early formation process. If we accepted that Kerr BHs are common at all scales, we expect that more objects with HFQPO detection will follow the universal scaling relation in the future (e.g., Alston et al. 2014b).

There is still no consensus on the *specific* oscillatory mechanism of the high-frequency, twin-peak BH QPOs. Two fundamental facts seem to be well-established, however, (1) These oscillations are characteristic of strong gravity. This follows from the $\nu \sim 1/M$ scaling first reported for the BH binaries by Remillard et al. (2002) (see also the Remillard & McClintock 2006 review article) and later generalized by Abramowicz et al. (2004) for all astrophysical BHs, including supermassive and (still hypothetical) intermediate-mass BHs; (2) These oscillations are characteristic of a non-linear resonance. This follows from the fact that their frequencies have a universal 3:2 ratio, as first noticed, and stressed, by Abramowicz & Kluźniak (2001).

We thank the referee for very helpful comments that improved the manuscript significantly. We thank L. J. Gou and S. N. Zhang for reading the manuscript and giving helpful comments. This work is supported by the NSFC grant 11033007, the Strategic Priority Research Program The Emergence of Cosmological Structures of the Chinese Academy of Sciences, grant No. XDB09000000 and the XTP project XDA04060604, EP project XDA04061100.

REFERENCES

- Abramowicz, M. A., Bulik, T., Bursa, M., & Kluźniak, W. 2003, *A&A*, 404, L21
- Abramowicz, M. A., & Kluźniak, W. 2001, *A&A*, 374, L19
- Abramowicz, M. A., Kluźniak, W., McClintock, J. E., & Remillard, R. A. 2004, *ApJL*, 609, L63
- Abramowicz, M. A., & Liu, F. K. 2012, *A&A*, 548, 3
- Alston, W. N., Markeviciute, J., Kara, E., Fabian, A. C., & Middleton, M. 2014a, *MNRAS*, 445, L16
- Alston, W. N., Parker, M. L., Markeviciute, J. et al. 2014b, *MNRAS*, preprint (arXiv:1411.0684)
- Aschenbach, B., Grosso, N., Porquet, D., & Predhl, P. 2004a, *A&A*, 417, 71
- Aschenbach, B. 2004b, *A&A*, 425, 1075
- Bian, W. H., & Huang, K. 2010, *MNRAS*, 401, 507
- Bloom, J. S., Giannios, D., Metzger, B. D., et al. 2011, *Sci*, 333, 203
- Blum, J. L., Miller, J. M., Fabian, A. C. et al. 2009, *ApJ*, 706, 60
- Burrows, D. N., Kennea, J. A., Ghisellini, G. et al. 2011, *Natur*, 476, 421
- Caballero-García, M. D., Belloni, T., & Zampieri, L. 2013, *MNRAS*, 436, 3262
- Dotti, M., Colpi, M., Pallini, S., Perego, A., & Volonteri, M. 2013, *ApJ*, 762, 68
- Feng, H., & Kaaret, P. 2007, *ApJ*, 668, 941
- Genzel, R., Schödel, R., Ott, T. et al. 2003, *Natur*, 425, 934
- Gierliński, M., Middleton, M., Ward, M., & Done, C. 2008, *Natur*, 455, 369
- Gillessen, S., Eisenhauer, F., Trippe, S. et al. 2009, *ApJ*, 692, 1075
- Gillessen, S., Genzel, R., Fritz, T. K. et al. 2012, *Natur*, 481, 51
- Homan, J., Miller, J. M., Wijnands, R. et al. 2005, *ApJ*, 623, 383
- Huang, C. Y., Wang, D. X., Wang, J. Z., & Wang, Z. Y. 2013, *RAA*, 13, 705
- King, A. R., & Kolb, U. 1999, *MNRAS*, 305, 654
- Liu, D., Pe'er, A., & Loeb, A. 2012, *ApJ*, preprint (arXiv:1211.5120)
- Liu, Z., Yuan, W., Lu, Y. J., & Zhou, X. L. 2014, *MNRAS*, preprint (arXiv:1411.4492)
- McClintock, J. E., Shafee, R., Narayan, R. et al. 2006 *ApJ*, 652, 518
- McClintock, J. E. et al. 2011, *CQGra*, 28, 4009
- Miller, J. M., & Gültekin, K. 2011, *ApJL*, 738, L13
- Miyoshi, M., Oyama, T., Shen, Z. et al. 2011, *ASPC*, 439, 336
- Pasham, D. R., Strohmayer, T. E., & Mushotzky, R. F. 2014, *Nature*, 513, 74
- Reid, M. J., McClintock, J. E., Steiner, J. F. et al. 2014, *ApJ*, 796, 2
- Remillard, R. A., Munro, M. P., McClintock, J. E., & Orosz, J. A. 2002, *ApJ*, 580, 1030
- Remillard, R. A. 2004, *AIPC*, 714, 13
- Remillard, R. A., & McClintock, J. E. 2006, *ARA&A*, 44, 49
- Reis, R. C., Fabian, A. C., Ross, R. R., Miller, J. M. 2009, *MNRAS*, 395, 1257
- Reis, R. C., Miller, J. M., Reynolds, M. T. et al. 2012, *Sci*, 337, 949
- Reynolds, C. S. 2014, *SSRv*, 183, 277
- Rezzolla, L., Yoshida, S., Maccarone, T. J., & Zanotti, O. 2003, *MNRAS*, 344, L37
- Shao, L., Zhang, F. W., Fan, Y. Z., & Wei, D. M. 2011, *ApJ*, 734, L33
- Shen, R. F., & Matzner, C. D. 2014, *ApJ*, 784, 87

- Steiner, J. F., Reis, R. C., McClintock, J. E. et al. 2011, MNRAS, 416, 941
- Strohmayer, T. E. 2001a, ApJL, 552, L49
- Strohmayer, T. E. 2001b, ApJL, 554, L169
- Strohmayer, T. E., & Mushotzky, R. F. 2003, ApJL, 586, L61
- Strohmayer, T. E., Mushotzky, R. F., Winter, L. et al. 2007, ApJ, 660, 580
- Titarchuk, L. 2002, ApJ, 578, L71
- Török, G., Abramowicz, M. A., Kluźniak, W., & Stuchlík, Z. 2005, A&A, 436, 1
- Török, G. 2005, A&A, 440, 1
- Volonteri, M., Madau, P., Quataert, E., & Rees, M. J. 2005, ApJ, 620, 69
- Wang, J. M., Hu, C., Li, Y. R., et al. 2009, ApJL, 697, L141
- Wang, J. Z., Lei, W. H., Wang, D. X., Zou, Y. C., Zhang, B., Gao, H., & Huang, C. Y. 2014, ApJ, 788, 32
- Zhang, S. N. 2013, FrPhy, 8, 630
- Zhou, X. L., Zhang, S. N., Wang, D. X., & Zhu, L. 2010, ApJ, 710, 16




FIG. 1.— Updated version of Figure 1 from Abramowicz et al. (2004); see also Figure 2 in Török (2005). In particular, scales and lines are similar and have the same meaning. Here, we merely add more data points which in 2004 were not available. Relationship between the quasi-periodic oscillation (QPO) frequency and the black hole mass. Three X-ray binaries, XTE J1550–564, GRO J1655–40, and GRS 1915+105, display a pair of high-frequency QPOs with a 3:2 frequency ratio. The frequencies of X-ray binaries are plotted for the stronger QPO that represent $2 \times \nu_0$. The solid line denotes the relation, ν (Hz) = $1862(M_{\text{BH}}/M_{\odot})^{-1}$ derived from three X-ray binaries (Remillard & McClintock 2006). The long-dashed line denotes the relation, ν (Hz) = $2030.8(M_{\text{BH}}/M_{\odot})^{-1}$ for a model of 3:2 resonance and spin parameter $a = 0.996$; the dotted-dashed line denotes the relation, ν (Hz) = $3068.9(M_{\text{BH}}/M_{\odot})^{-1}$ for a model of 3:1 resonance and spin parameter $a = 0.996$; the dotted line denotes the Kepler frequency for a non-spinning Schwarzschild black hole at the innermost stable circular orbit. The QPO reported in the active galactic nucleus REJ 1034+396, the ultra-luminous X-ray source NGC 5408 X-1, and the tidal disruption event Swift J1644+57 are plotted. The observed QPO frequencies in the ultra-luminous X-ray source M82 X-1 display 3:2 ratio (Pasham et al. 2014). The frequency for this object also represents $2 \times \nu_0$. The Galactic Center Sgr A* follows the 3:1 resonance line, although the significance of QPOs in Sgr A* is still under debate. See the text for details.

TABLE 1
LISTS OF OBJECTS WITH (HF)QPO DETECTION

Source Name	frequency (Hz)	$2\nu_0$	Ref1.	M_{BH} ($M_{\odot}$)	Method	Ref2.
GRO J1655 – 40	300 ± 10	Yes	1	6.3 ± 0.3	D	9
XTE J1550 – 564	180 ± 8	Yes	1	9.6 ± 1.2	D	9
GRS 1915 + 105	113 ± 5	Yes	2	$12.4^{+2.0}_{-1.8}$	D	10
H 1743 – 322	163^{+5}_{-2}	Yes	3	...	...	...
M82 X–1	3.32 ± 0.06	Yes	4	415 ± 63	P	4
NGC 5408 X–1	$(20.2 \pm 0.3)\text{E-3}$	No	5	$(1 \pm 0.3)\text{E5}$	T	11
Swift J1644 + 57	$(4.8 \pm 0.3)\text{E-3}$	No	6	$(3.16^{+35.6}_{-2.9})\text{E5}$	R	12
RE J1034 + 396	$(2.68 \pm 0.1)\text{E-4}$	No	7	$(4^{+3}_{-1.5})\text{E6}$	T	13
Sgr A*	$(1.07 \pm 0.05)\text{E-3}$	Yes	8	$(4.31 \pm 0.42)\text{E6}$	D	14

Ref1. –1: Remillard et al. (2002); 2: Remillard (2004); 3: Homan et al. (2005); 4: Pasham et al. (2014); 5: Strohmayer et al. (2007); 6: Reis et al. (2012); 7: Gierliński et al. (2008); 8: Aschenbach et al. (2004a). Ref2. –9: Remillard & McClintock (2006); 10: Reid et al. (2014); 11: Huang et al. (2013); 12: Miller & Gültekin (2011); 13: Zhou et al. (2010); 14: Gillessen et al. (2009). Method for black hole mass estimates. –D: Dynamical method; P: relativistic precession model; T: X-ray timing; R: radio observations.